arXiv:1209.5327v1 [quant-ph] 24 Sep 2012

Non-adiabatic control of quantum energy transport in ordered arrays

Ping Xiang, Marina Litinskaya, Evgeny A. Shapiro, and Roman V. Krems[1]

[1] *Department of Chemistry, University of British Columbia, Vancouver, V6T 1Z1, Canada*

An elementary excitation in an aggregate of coupled particles generates a collective excited state. We show that the dynamics of these excitations can be controlled by applying an instantaneous external potential that modifies the quantum phase of individual particles. We show that specific phase transformations can be used to accelerate or decelerate quantum energy transport and spatially focus delocalized excitations onto different parts of ordered arrays of quantum particles.

PACS numbers: 42.50.Hz, 34.50.Ez, 42.50.Dv, 37.10.Pq

Many physical systems, such as molecular dye aggregates[1], photosynthetic complexes [2], atoms and molecules in optical lattice traps [3, 4] or self-assembled monolayers on solid substrates [5], represent a chain of coupled monomers, which, once excited, can exchange the excitation energy. This leads to energy transport. Understanding and controlling energy transport is crucial for many applications such as building efficient light-harvesting devices [6] and scalable quantum computing networks [7]. In this work, we explore the general algorithm for controlling transport through an array of coupled quantum monomers by applying monomer-specific external perturbations. This is possible to realize in an ensemble of spatially separated monomers, such as Rydberg atoms or molecules trapped on an optical lattice.

Excitation of a coupled many-body system generates a wavepacket representing a coherent superposition of single-site excitations. We explore the possibility of using external fields to control their dynamics. The method proposed here is based on shaping the many-body wavepackets by a series of sudden perturbations, in analogy with the techniques developed for strong-field alignment and orientation of molecules in the gas phase [8]. Alignment is used in molecular imaging experiments and molecular optics [8, 9], and is predicted to provide control over mechanical properties of molecular scattering [10]. Our technique allows for the possibility to study condensed-matter excitations and energy transport without statistical averaging. We demonstrate that a delocalized excitation can be spatially focussed, enabling directed transport of energy to a particular monomer. This can be exploited for the experimental realization of controlled chemical transformations and as a toolbox for optimizing photon-harvesting systems.

We consider an ensemble of N coupled identical monomers possessing two internal states $|g\rangle$ and $|e\rangle$ arranged in a one-dimensional array with translational symmetry. The excitation spectrum of the system is determined by two parameters: the monomer excitation energy ΔE_{e-g} and the coupling strength $\alpha(n)$ between the monomers separated by the distance na, where a is the nearest neighbor separation and n is an integer. The singly excited state of the system is

$$|\psi_{\rm exc}\rangle = \sum_{n=1}^{N} C_n |e_n\rangle \prod_{i\neq n} |g_i\rangle, \qquad (1)$$

where $|g_n\rangle$ and $|e_n\rangle$ denote the ground and excited states in site n, $C_n = e^{iakn}/\sqrt{N}$ and $|\psi_{\rm exc}\rangle \Rightarrow |\psi_{\rm exc}(k)\rangle$ represents a quasi-particle called Frenkel exciton, characterized by the wave vector k [11]. The energy of the exciton is given by $E(k) = \Delta E_{e-g} + \alpha(k)$ with $\alpha(k) = \sum_n \alpha(n) e^{-iakn}$. In the nearest neighbor approximation,

$$E(k) = \Delta E_{e-g} + 2\alpha \cos ak, \qquad (2)$$

where $\alpha = \alpha(1)$. A localized excitation placed on a single site (or a limited number of $M < N$ sites) is a coherent superposition of the exciton states $|\psi_{\rm exc}(k)\rangle$ with different k.

Control over energy transport in an ordered array can be achieved by (i) shifting the exciton wavepackets in the momentum representation (which modifies the group velocity and the shape evolution of the wavepackets) and (ii) focusing the wavepackets in the coordinate representation to produce localized excitations in an arbitrary part of the lattice. To achieve this, we propose to apply a series of site-dependent perturbations. For (i), the essential idea is to add a factor $e^{i\delta an}$ to each term in the expansion (1), so that each $|\psi_{\rm exc}(k)\rangle$ is transformed into $|\psi_{\rm exc}(k+\delta)\rangle$. This transformation shifts the wavepackets by δ in k-space while preserving their shape. As a result, one can engineer wavepackets probing any part of the dispersion $E(k)$ leading to different group velocity and shape evolution.

Adding a site-dependent phase to the excitonic wavefunction exploits an interplay of the adiabatic and sudden time scales. Consider, first, the n-th monomer subjected to an external field $\mathcal{E}_n(t)$ which varies from 0 to some value and then back to 0 in time T. If the variation is adiabatic with respect to the evolution of the free monomer states, $T \gg \hbar/\Delta E_{e-g}$, each eigenstate $|f\rangle$ of the monomer acquires a state-dependent phase shift [12]

$$|f_n(T)\rangle = e^{-i\phi_n^f} |f_n(0)\rangle, \text{ where } \phi_n^f = \frac{1}{\hbar}\int_0^T E_n^f(t)dt, \qquad (3)$$

$E_n^f(t)$ is the instantaneous eigenenergy and f can be e or g. Now consider the action of such phase change on the collective excitation state (1). If $T \ll \hbar/\alpha$, the change is sudden with respect to the excitation transfer between monomers and the state (1) acquires a site-dependent phase $\Phi_n = \phi_n^e - \phi_n^g$. If $\Phi_n = \Phi_0 + na\delta$, then the momentum δ is imparted onto the excitonic wavefunction. By analogy with "pulsed alignment of molecules" [8], we call this transformation a "phase kick" or "momentum kick". Its action is also similar to that of a thin prism on a wavefront of a monochromatic laser beam.

In order to illustrate the shifting of exciton wavepackets in the momentum space, we solve numerically the time-dependent Schrödinger equation with the unperturbed Hamiltonian $H_{\rm exc} = \Delta E_{e-g} \sum_n |e_n\rangle\langle e_n| + \sum_{n,m} \alpha(n-m)|e_n, g_m\rangle\langle g_n, e_m|$, subjected to a transient site-dependent external perturbation. The excitation at $t = 0$ is described by a Gaussian wave packet of the exciton states $|\psi_{\rm exc}(k)\rangle$, with the central wavevector $k = 0$. The external perturbation is provided by a spatially inhomogeneous pulse of electromagnetic field. Fig. 1 shows that the entire wave packet acquires momentum during the pulse (left panels). This is manifested as a phase variation in the coordinate representation, and as a shift of the central momentum in the k-representation. After the external perturbation is gone, the wave packet does not evolve in the k-representation and moves with the acquired uniform velocity in the coordinate representation.

In optics, a thin lens focuses a collimated light beam by shifting the phase of the wavefront, thus converting a plane wave to a converging spherical wave. Similarly, a phase kick can serve as a time domain "lens" for collective excitations: an excitation delocalized over a large number of monomers can be focused onto a small region of the array. By analogy with optics, a concave symmetric site-dependent phase $\Phi(n)$ applied simultaneously to all monomers may turn a broad initial distribution $C_n(t = 0)$ into a narrow one. The dynamics of the amplitudes in Eq. (1) is given by

$$C_m(t) = \frac{1}{N} \sum_{n,k} C_n(t=0) e^{i[\Phi(n) + ka(m-n) - E(k)t/\hbar]} . \quad (4)$$

The shape of the dispersion curve $E(k)$ is crucial for the focusing efficiency: while waves with quadratic dispersion can be focused, those with linear dispersion propagate without changing the wavepacket shape [13]. The cosine-like dispersion of excitons (2) offers both the quadratic (at low k) and linear (at $k \approx \pm\pi/2a$) regions. A rigorous analysis shows that, for the cosine dispersion, the wave packet (1) with $C_n = J_{n-n_0}(2\alpha\tau)e^{i\pi(n-n_0)/2}$, where J_n is the Bessel function of order n, must focus onto the single site n_0 at time τ. However, creating such a wavepacket may require multiple and complicated phase transformations. For practical applications, it may be easier to constrain all phase transformations such that the wavepacket always remains in the quadratic part of the cosine dispersion (2). For such wavepackets, adding a quadratic phase $\Phi(n) = \Phi_0(n - n_0)^2$ leads to focusing around site n_0. Below we illustrate the effect of the quadratic phase transformation for two types of initial states.



FIG. 1: (Color online) Evolution of the exciton wavepacket in the k and real spaces. The phase of the wave function is shown by color. The calculation is for a one-dimensional array of 201 monomers with $\alpha = 22.83$ kHz and $\Delta E_{e-g} = 12.14$ GHz, which corresponds to LiCs molecules trapped on an optical lattice with $a = 400$ nm and subjected to a homogenous DC field of 1 kV/cm directed perpendicular to the intermolecular axis. The kicking potential is provided by a $\lambda = 1024$ nm Gaussian laser beam, with the propagation direction along the array axis, focused to 5 μm, with the intensity at the focus equal to 10^7 W/cm^2. The laser pulse is on between 0 and 3 μs. The molecules are placed on the beam axis with the first molecule 5 μm away from the focus. Thus, the laser intensity is gradually decreased along the array.

First, consider a broad Gaussian wavepacket (1) with $C_n(\sigma_x; t = 0) = \sqrt{a/\sigma_x\sqrt{\pi}} \exp\left[-a^2(n-n_0)^2/2\sigma_x^2\right]$ where $\sigma_x \gg a$. By applying the quadratic phase to the Gaussian wave packet with quadratic dispersion, one creates a wave packet with the width in the k-representation [13]

$$\sigma_k(\sigma_x, \Phi_0) = \frac{1}{\sigma_x}\sqrt{1 + 4\Phi_0^2\sigma_x^4/a^4}. \quad (5)$$

Eq. (5) shows that the value of Φ_0 determines the degree of focusing. However, for systems with dispersion (2), large values of Φ_0 may take the wavepacket outside the quadratic part of the dispersion. To find the optimal phase Φ_0^* that keeps the wavepacket within the quadratic dispersion while focusing it, we use the condition $\Delta_k = a\sigma_k \lesssim 1$, which yields $\Phi_0^* = a/2\sigma_x$ for the optimal focusing. At time $t_* \approx 1/4\alpha\Phi_0^*$, the wavepacket is most focused and has a width

$$\sigma_{x,F}(\Phi_0^*) = \frac{\sigma_x}{\sqrt{1 + 4{\Phi_0^*}^2\sigma_x^4/a^4}} \approx a. \quad (6)$$

Second, consider a completely delocalized excitation (1) with $C_n(k;t=0) = e^{iakn}/\sqrt{N}$ describing an eigenstate of an ideal system of N coupled monomers. If $E(k)$ in Eq. (4) is assumed to be $E(k) = \Delta E_{e-g} - \alpha a^2 k^2$, the phase $\Phi(n) = \Phi_0(n-n_0)^2$ with $\Phi_0 = \pi/N$ reduces the amplitudes C_m at $t_* = N/4\pi$ to $C_m(k;t_*) \approx \sqrt{i}e^{-iNa^2k^2/4\pi}\delta_{n,n_0-\nu_k}$, where $\nu_k = Nak/2\pi$ [14]. However, an estimate shows [14] that this phase modulation would create a wavepacket with the width $\Delta_k(\Phi_0) \approx 2N\Phi_0$ in the momentum space spanning the entire range of k-values. For an arbitrary value of $\Delta_k(\Phi_0)$, the site amplitudes at the time of focusing $t_* = 1/4\alpha\Phi_0$ are [14]

$$C_n(k;t=t_*) \approx \frac{e^{i\Delta(\Phi_0)n^2/2N}}{n}\sqrt{\frac{2i}{\pi\Delta_k(\Phi_0)}}\sin(n\Delta_k(\Phi_0)/2). \tag{7}$$

In order to keep the linear part of the dispersion spectrum unpopulated, we choose $\Phi_0^* \sim 1/2N$ so that $\Delta_k(\Phi_0^*) \sim 1$.



FIG. 2: Focusing of a completely delocalized collective excitation (panels a and b) and a broad Gaussian wave packet of Frenkel excitons (panels c and d) using the quadratic phase transformations at $t = 0$ as described in text. The dashed lines show the initial distribution magnified by 20 and 5 respectively in (b) and (d). The solid curves in panels (b) and (d) correspond to two different phase transformation focusing the same wave packet onto different parts of the array. The calculations are performed with the same parameters α, a, and ΔE_{e-g} as in Figure 1. The results in panels (a) and (b) are computed with nearest and second neighbor couplings only; the results in panels (c) and (d) are computed with all couplings accounted for.

Eqs. (6) and (7) are valid for a many-body system with nearest neighbor interactions only. In most physical systems, the energy dispersion is modified by long-range couplings. In order to confirm that the above predictions are also valid for systems with long-range interactions and illustrate the focusing of delocalized excitations, we compute the time evolution of the wave packets by solving the wave equation numerically for a system with long-range dipole - dipole interactions. Figure 2 illustrates the focusing dynamics of a completely delocalized excitation in a system with nearest neighbor and second-order couplings (panels a and b) and a broad Gaussian wave packet in a system with all long-range couplings explicitly included in the calculation (panels c and d). The results show that, although the long-range couplings lead to an oscillating interference pattern, the collective excitations can be focussed to a few lattice sites.

By analogy with optics, it should be expected that the focusing phase transformations must be more effective in systems of higher dimensionality. To illustrate this, we repeated the calculations presented in Figures 2c and 2d for a delocalized excitation placed in a 2D array with an external potential that modulates the phase as a function of both x and y. Figure 3 shows the focusing of an initially broad wave packet onto different parts of a 2D lattice induced by the quadratic phase transformation. The calculations include all long-range couplings. Further, similar to spatial light modulators in optics, multiple phase kicks of varying profiles separated by periods of free evolution may enable arbitrary shaping of the excitonic wave packet in the coordinate and k-spaces.



FIG. 3: Focusing of a delocalized excitation in a 2D array shown at $t = 0$ in panel (a) onto different parts of the lattice (panels b–d). The probability distribution in panel (a) is enhanced by the factor 72 for clarity. The calculations are performed with the same parameters α, a, and ΔE_{e-g} as in Figure 1 and the quadratic phase transformation at $t = 0$.

While the results shown in Figures 1-3 are general, for concreteness, the calculations presented are performed for an experimentally realizable system of polar diatomic molecules in the $^1\Sigma$ electronic state trapped on an optical lattice [15]. We assume that the optical lattice is prepared with one molecule per lattice site and that tunneling of molecules between different lattice sites is suppressed. The rotational states of the molecules are labeled by the quantum number of the rotational angular momentum $\boldsymbol{J}$ and the projection M_J of $\boldsymbol{J}$ on the space-fixed quantization axis Z. We choose the rotational ground state $|J = 0, M_J = 0\rangle$ as $|g\rangle$ and the rotational excited state $|J = 1, M_J = 0\rangle$ as $|e\rangle$. The state $|J = 1, M_J = 0\rangle$ is degenerate with the states $|J = 1, M_J = \pm 1\rangle$. This degeneracy can be lifted by applying a homogeneous DC electric field, making the $|g\rangle$ and $|e\rangle$ states an isolated two-level system. The molecules in different lattice sites are coupled by the dipole - dipole interaction $V_{\rm dd}(n-m)$. The separation between adjacent sites of an optical lattice is equal to half the wavelength of the trapping laser and is, typically, in the range between 256 and 800 nm [4]. The magnitude of the coupling constant $\alpha(n-m) = \langle e_n, g_m|V_{\rm dd}(n-m)|g_n, e_m\rangle$ between molecules with the dipole moment 1 Debye separated by 500 nm is on the order of 1 kHz [16].

For molecules on an optical lattice, one can implement the phase kicks by modifying the molecular energy levels with pulsed AC or DC electric fields. The rotational energy levels for $^1\Sigma$ molecules in a combination of weak AC and DC electric fields are given by [17]

$$E_{J,M_J} \approx BJ(J+1) + \frac{\mu^2\mathcal{E}_{DC}^2}{2B}G(J,M_J) - \frac{\alpha_\perp\mathcal{E}_{AC}^2}{4} + \frac{(\alpha_{||}-\alpha_\perp)\mathcal{E}_{AC}^2}{4}F(J,M_J) \quad (8)$$

where B is the rotational constant, $G(0,0) = -1/3$, $G(1,0) = 1/5$, $F(0,0) = 1/3$, $G(1,0) = -1/5$, $\mathcal{E}_{AC}$ is the envelope of the quickly oscillating AC field, $\alpha_\parallel$ and $\alpha_\perp$ are the parallel and perpendicular polarizabilities and μ is the permanent dipole moment of the molecule.

The momentum shift of the exciton wave packets can be achieved by applying a time-varying DC electric field $\mathcal{E}(t) = \mathcal{E}_* + \mathcal{E}(n)\sin^2(\pi t/T)$, where $\mathcal{E}(n)$ is linear with respect to n. Assuming that $\mathcal{E}(n) = (n - n_0)A$ and $\mathcal{E}(n) \ll \mathcal{E}_*$, and using Eqs. (3) and (8), gives $\delta = 4A\mathcal{E}_*\mu^2T/15\hbar Ba$. We have confirmed this result by a numerical computation showing that for LiCs molecules in an electric field of $\mathcal{E}_* = 1$ kV/cm, an electric field pulse with $A = 7.434 \times 10^{-4}$ kV/cm and $T = 1$ μs results in a kick of $\delta = \pi/2a$, bringing an excitonic wave packet from the $k = 0$ region to the middle of the dispersion zone.

An alternative strategy is to use a pulse of an off-resonance laser field. Consider a laser pulse whose intensity varies linearly across the molecular array:

$$\mathcal{E}_{AC,n}^2(t) = [\mathcal{E}_0^2 + (n - n_0)\mathcal{E}_1^2]\,\sin^2(\pi t/T) \quad (0 < t < T)\ . \quad (9)$$

Using Eqs. (3) and (8), we estimate the momentum kick by such a pulse as $\delta = T\mathcal{E}_1^2(\alpha_\parallel - \alpha_\perp)/15a$. The results shown in Figure 1 correspond to this set of parameters.

In addition to the momentum kicking technique, energy transport in an array of molecules on an optical lattice can be controlled also by varying the angle θ between the intermolecular axis and the applied DC field. This angle influences the dipole-dipole interaction between the molecules and determines the sign and magnitude of α, and therefore the shape of the dispersion curve, as shown in Figure 4(a). This enables control over the sign and magnitude of the group velocity of an excitonic wave packet with $k \neq 0$. Dynamically tuning θ, one can propagate a localized excitation to different parts of the lattice, as shown in Figure 4b.



FIG. 4: (Color online) (a) Exciton dispersion curves for a 1D ensemble of diatomic molecules on an optical lattice for different angles θ between the direction of the external DC electric field and the axis of the molecular array; (b) Propagation of a wave packet centered at $ak = -1.0464$ controlled by tuning the electric field direction. Thin dashed line depicts the corresponding angle variations with time.

In summary, any elementary excitation in an aggregate of coupled monomers can be represented as a coherent superposition of free exciton states. We propose a method for shaping the exciton wave packets using “phase kicking” perturbations that temporarily modulate the energy levels of the monomers leading to a displacement of the wave packets in the momentum representation. This can be used to control the energy transport in ordered ensembles of quantum particles. In particular, we show that the phase kicking technique can be used for focusing a delocalized collective excitation to a particular part of a 1D or 2D lattice.

Control over excitation transport is needed for creating networks of quantum processors where information is transmitted over large distances with photons and stored in arrays of quantum monomers via one of the quantum memory protocols [18]. Momentum kicking can be used for information transport within a single array. Focusing excitonic wave packets enables local storage of information, while controlled interactions of multiple excitons [19] or excitons with lattice impurities [20] may be used to implement logic gates.

Controlled energy transport in molecular arrays may also be used for the study of controlled chemical interactions for a class of reactions stimulated by energy excitation of the reactants. Directing quantum energy to a particular lattice site containing two or more reagents can be used to induce a chemical interaction [21], an inelastic collision or predissociation [22] with the complete temporal and spatial control over the reaction process.

Acknowledgment. This work was supported by NSERC of Canada and the Peter Wall Institute for Advanced Studies.

References

---

[1] F. Würthner, T. E. Kaiser, C. R. Saha-Möller, Angew. Chem. Int. Ed. **50**, 3376 (2011)

[2] G. S. Engel, T. R. Calhoun, E. L. Read, Tae-Kyu Ahn, T. Mančal, Yuan-Chung Cheng, R. E. Blankenship, and G. R. Fleming, Nature **446**, 782 (2007), and references therein

[3] I. Bloch, Nat. Phys. **1**, 23 (2005)

[4] G. Grynberg and C. Robilliard, Phys. Rep. **355** 335 (2001); R. Grimm, M. Weidemüller, and Yu.B. Ovchinnikov, Adv. At. Mol. Opt. Phys. **42**, 95 (2000)

[5] J. C. Love, L. A. Estroff, J. K. Kriebel, R. G. Nuzzo, and G. M. Whitesides, Chem. Rev. **105**, 1103 (2005)

[6] B. A. Gregg, J. Phys. Chem. B **107**, 4688 (2003)

[7] D. P. DiVincenzo, Fort. Phys. **48**, 771 (2000)

[8] T. Seideman and E. Hamilton, Adv. At. Mol. Opt. Phys. **52**, 289 (2006) and references therein;

[9] I. V. Litvinyuk, K. F. Lee, P. W. Dooley, *et. al.* Phys. Rev. Lett. **90**, 233003 (2003); S. M. Purcell and P. F. Barker , Phys. Rev. Lett. **103**, 153001 (2009); E. Gershnabel, M. Shapiro and I. Sh. Averbukh,J. Chem. Phys. **135**, 194310 (2011).

[10] U. Steinitz, Y. Prior, I. Sh. Averbukh, Phys. Rev. Lett. **109** 033001 (2012); Yu. Khodorkovsky, J. R. Manson, I. Sh. Averbukh, Phys. Rev. A **84**, 053420 (2011).

[11] V. Agranovich, Excitations in Organic Solids (Oxford University Press, Oxford, 2008)

[12] M. H. S. Amin, Phys. Rev. Lett. **102**, 220401 (2009), and references therein

[13] M. V. Fedorov, Atomic and Free Electrons in a Strong Light Field (World Scientific, Singapore, 1997); J. W. Goodman, Introduction to Fourier Optics (2nd ed.) (McGraw-Hill, New York, 1996)

[14] We use the approximate equality

$$\int_{-M}^{M} dx\, e^{-i(ax^2+bx)} \approx \sqrt{\frac{\pi}{ia}}\, e^{ib^2/4a}\, \Theta(-2Ma < b < 2Ma),$$

where $\Theta(z) = 1$ if z is true and zero otherwise. It was derived by approximating the error function of a complex argument $\mathrm{Erf}(\sqrt{i}x)$ by the sign function – the approach which works the better the larger x.

[15] A. Chotia, B. Neyenhuis, S. A. Moses, B. Yan, J. P. Covey, M. Foss-Feig, A. M. Rey, D. S. Jin, and J. Ye, Phys. Rev. Lett. **108**, 080405 (2012).

[16] F. Herrera and R. V. Krems, Phys. Rev. A **84**, 051401(R) (2011)

[17] B. Friedrich and D. Herschbach, J. Phys. Chem. **99**, 15686 (1995).

[18] A.I. Lvovsky, B.C. Sanders and W. Tittel, Nature Photonics **3**, 706 (2009), and references therein

[19] P. Xiang, M. Litinskaya, and R. V. Krems, Phys. Rev. A **85**, 061401(R) (2012)

[20] A. J. Daley, S. R. Clark, D. Jaksch, P. Zoller, Phys. Rev. A **72**, 043618 (2005)

[21] R. V. Krems, Physical Chemistry Chemical Physics **10**, 4079 (2008)

[22] A. O. G. Wallis and R. V. Krems, J. Chem Phys **135**, 124313 (2011)